

A consensus genetic map of cowpea [*Vigna unguiculata* (L) Walp.] and synteny based on EST-derived SNPs

Wellington Muchero^{a,1}, Ndeye N. Diop^{b,1}, Prasanna R. Bhat^{b,2}, Raymond D. Fenton^b, Steve Wanamaker^b, Marti Pottorff^b, Sarah Hearne^c, Ndiaga Cisse^d, Christian Fatokun^c, Jeffrey D. Ehlers^b, Philip A. Roberts^a, and Timothy J. Close^{b,3}

Departments of ^aNematology and ^bBotany and Plant Sciences, University of California, Riverside, CA 92521; ^cThe International Institute of Tropical Agriculture, Oyo Road, PMB 5320, Ibadan, Nigeria; and ^dSenegalese Institute of Agricultural Research, Bambey, Senegal

Edited by Deborah P. Delmer, Rockefeller Foundation, New York, NY, and approved September 4, 2009 (received for review May 30, 2009)

Consensus genetic linkage maps provide a genomic framework for quantitative trait loci identification, map-based cloning, assessment of genetic diversity, association mapping, and applied breeding in marker-assisted selection schemes. Among “orphan crops” with limited genomic resources such as cowpea [*Vigna unguiculata* (L.) Walp.] ($2n = 2x = 22$), the use of transcript-derived SNPs in genetic maps provides opportunities for automated genotyping and estimation of genome structure based on synteny analysis. Here, we report the development and validation of a high-throughput EST-derived SNP assay for cowpea, its application in consensus map building, and determination of synteny to reference genomes. SNP mining from 183,118 ESTs sequenced from 17 cDNA libraries yielded $\approx 10,000$ high-confidence SNPs from which an Illumina 1,536-SNP GoldenGate genotyping array was developed and applied to 741 recombinant inbred lines from six mapping populations. Approximately 90% of the SNPs were technically successful, providing 1,375 dependable markers. Of these, 928 were incorporated into a consensus genetic map spanning 680 cM with 11 linkage groups and an average marker distance of 0.73 cM. Comparison of this cowpea genetic map to reference legumes, soybean (*Glycine max*) and *Medicago truncatula*, revealed extensive macrosynteny encompassing 85 and 82%, respectively, of the cowpea map. Regions of soybean genome duplication were evident relative to the simpler diploid cowpea. Comparison with *Arabidopsis* revealed extensive genomic rearrangement with some conserved microsynteny. These results support evolutionary closeness between cowpea and soybean and identify regions for synteny-based functional genomics studies in legumes.

legume | diploid | recombinant inbred lines | segregating traits | 1536-plex genotyping

Recent progress in genome resource development for model and major crop plants has energized genetic research and fostered a surge of new initiatives in plant improvement. However, this activity has largely bypassed “orphan crops” such as cowpea [*Vigna unguiculata* (L.) Walp.] ($2n = 2x = 22$), which are crops of relevance to food security and income for subsistence farmers in developing countries (1). Despite the limited genome resources, access to most of the genes in these organisms can be gained through cDNA sequences, which represent expressed genes. Partial cDNA sequences, known as ESTs, when determined from multiple genotypes of a species, facilitate the identification of SNPs in protein-encoding genes and can be used in conjunction with mapping populations to generate genetic linkage maps that represent a gene-based framework of the genome.

Cowpea is a very important leguminous crop of the developing world. The crop is particularly important in sub-Saharan Africa, where >10 million hectares are cultivated in the semiarid Savanna and Sahelian zones of West and Central Africa (<http://faostat.fao.org>). Several parts of South America (particularly north-eastern Brazil and Peru) and parts of south Asia (India, Myanmar), the Middle East, and the southern regions of North America are also important cowpea production regions (2). Cowpea is a

particularly valuable component of low-input farming systems of resource-poor farmers because of its productivity and yield stability in the face of abiotic stress (drought, heat, low soil fertility), and the ability of the crop to enhance soil fertility for succeeding cereal or tuber crops grown in rotation (3). With its greater tolerance to heat, drought, and low soil fertility (4) and yet close evolutionary relatedness to other economically important grain legumes such as common bean (*Phaseolus vulgaris*) and soybean (*Glycine max*), cowpea can serve as a model species for crop adaptation to these stresses. Until recently only limited progress had been made in basic gene discovery and gene regulation in cowpea, with such information available for *Vigna* species being 2- to 5-fold less than for pea (*Pisum sativum*), common bean (*Phaseolus vulgaris*), and alfalfa (*Medicago sativum*) and >500 -fold less than for soybean (*G. max*) (5).

Before this work, only cross-specific and low-density genetic linkage maps comprised mostly of anonymous markers were available for cowpea. To date, the most comprehensive genetic map of *V. unguiculata* consists of 11 linkage groups (LGs) spanning a total of 2,670 cM, with an average distance of ≈ 6 cM between markers. It includes 242 amplified fragment length polymorphism (AFLP) and 18 disease and pest resistance-related markers (6), plus 133 random amplified polymorphic DNA (RAPD), 39 restriction fragment length polymorphism (RFLP), and 25 AFLP markers from an earlier map (7), for a total of 441 markers. However, the cM distance covered by this map is three to four times greater than other published RFLP-, RAPD-, and AFLP-based cowpea linkage maps (7–9).

Here, we describe the development and implementation of high-throughput SNP genotyping in cowpea and its application to produce a high-density SNP consensus map based on genotyping 741 members of six recombinant inbred line (RIL) populations, which can be related readily to prior maps through shared markers. Synteny was investigated to *G. max*, a major crop of worldwide importance, and which outside of the genus *Vigna* and next to common bean (*P. vulgaris*) is the closest legume relative of cowpea (10). Synteny with *M. truncatula* and *Arabidopsis thaliana* was also surveyed.

Results

Identification of SNPs and Development of GoldenGate Assays. Sequencing 11 cDNA libraries generated 141,453 ESTs (Table S1). In

Author contributions: J.D.E., P.A.R., and T.J.C. designed research; W.M., N.N.D., P.R.B., R.D.F., M.P., J.D.E., and T.J.C. performed research; S.H., N.C., C.F., and J.D.E. contributed new reagents/analytic tools; W.M., N.N.D., P.R.B., R.D.F., S.W., M.P., J.D.E., P.A.R., and T.J.C. analyzed data; and W.M., N.N.D., J.D.E., P.A.R., and T.J.C. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

¹W.M. and N.N.D. contributed equally to this work.

²Present address: Monsanto Research Centre, Bangalore, India, 560092.

³To whom correspondence should be addressed. E-mail: timothy.close@ucr.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0905886106/DCSupplemental.

Table 1. Summary of genotypes used for SNP discovery

Genotype	Other identifier	Origin	Type	Key traits
CB27	H9–8–27	UCR	Cultivar, MP	Heat tolerance, <i>Fusarium</i> wilt & root-knot nematode resistance
IT845-2049	UCR 430	IITA	Breeding line MP	Root-knot nematodes resistance
524B		UCR	Breeding line, MP	Large seed, <i>Fusarium</i> wilt resistance
UCR 5301	TVNu-463	Botswana	Wild ssp. <i>dekindtiana</i>	Genetic diversity
UCR 779	Botswana 19A	Botswana	Landrace	Cowpea aphid resistance
IT97K-461–4		IITA	Breeding line	<i>Striga gesnerioides</i> resistance
UCR 707	Ex-Luanda	Kenya	Landrace	Genetic diversity
PI 418979		China	Landrace	Asparagus bean
UCR 41	TVu-1996	Nicaragua	Landrace	Genetic diversity
UCR 2563	TVu-1522	Iran	Landrace	Genetic diversity
Dan Ila		IITA	Breeding line, MP	Drought tolerance
TVu 11986		IITA	Breeding line, MP	Drought tolerance
TVu 7778		IITA	Breeding line, MP	Drought susceptible
12008D		ILRI	Landrace	Animal feed, drought tolerance
1393–2-1		UCR	Breeding line	Chilling tolerance, large seed

Three additional libraries contributed ESTs but were not used for SNPs because trace files were not available. MP, mapping parent for RILs used in this work or otherwise. University of California, Riverside; IITA, International Institute of Tropical Agriculture; ILRI, International Livestock Research Institute.

addition, 41,505 ESTs from two libraries produced in a Generation Challenge Program project led by the International Institute of Tropical Agriculture and 160 from other libraries were included to give a total of 183,118 ESTs, which were used to create a comprehensive EST sequence assembly. HarvEST:Cowpea (<http://harvest.ucr.edu>) provides this information, which is also summarized in [Table S1](#). The total number of nucleotides in the consensus sequences from contigs containing at least four members was 11.3 Mbp. Taking 10,000 as the number of high-quality SNPs, the average SNP frequency was 1 per 1.13 kbp.

The 1,536-SNP GoldenGate assay (see *Materials and Methods*) was applied to 759 DNA samples from mapping parents, RILs, and synthetic heterozygotes. Of these, 1,375 SNPs had satisfactory technical performance, an 89.55% success rate. Genotypes used for the 11 cDNA libraries (Table 1) contained no heterozygous loci.

Polymorphism and Segregation Distortion. A total of 986 of the 1,375 SNPs (72%) were polymorphic in at least one RIL population. In total, 410 markers exhibited segregation distortion in one or more populations; however, 360 of these had minor allele frequencies (MAFs) higher than the 0.30 threshold in at least one population and could be mapped. Of the remaining 50 with low MAF, 39 were polymorphic in only one population, and 11 were polymorphic in multiple populations but with low MAF in each case. SNPs with a low MAF (< 0.30) in each population ranged from 33 for RIL 524B \times IT84S-2049 to 236 for RIL Dan IIa \times TVu-7778 (Table 3). Six SNPs were discarded because they had GoldenGate Assay “no-call” frequencies $>5\%$ in the population in which they were segregating, and two SNPs were eliminated after they mapped in two different LGs.

A total of 58 SNPs, mainly from the 3×3 list (see *Materials and Methods*) or dependent on cowpea genotype UCR779 in the pairwise list, could not be mapped in any population. As a net result, 928 SNPs that had MAF at least 0.3 in one or more mapping populations were used in map construction. In pairwise comparisons, between 99 and 202 SNPs were shared by any two mapping populations. The numbers of SNPs shared between populations are summarized in Table 2.

Individual Genetic Linkage Maps. Elimination of RILs with nonparental alleles, excessive heterozygosity, and excessive no-calls (see *Materials and Methods*) resulted in the final mapping population sizes given in Table 3. Therefore, a total of 632 RILs were used in individual and consensus map building. For each of the six mapping populations, LGs were resolved without conflicting marker assignments by using JoinMap 3.0 (11) and the parameters described in *Materials and Methods*. The stringent mapping parameters (see *Materials and Methods*) adopted for individual map construction resulted in the number of LGs ranging from the expected 11 in TVu-14676 \times IT84S-2246-4 to 15 in 524B \times IT84S-2049. Although the additional smaller LGs could be consolidated into larger LGs using lower logarithm of odds (LOD) thresholds, this separation was tolerated before consensus mapping to minimize spurious linkage that could result in tangled consensus LGs. Maps of the six populations ranged from 600 to 665 cM and consisted of 288–436 SNP markers. The largest gap between mapped SNPs ranged from 13.3 to 17.9 cM for individual maps (Table 3). Average marker distances ranged from 2.31 cM in Dan Ila \times TVu-7778 to 1.52 cM in 524B \times IT84S-2049.

Consensus Genetic Linkage Map. Because of the high prevalence of segregation distortion in the Dan IIa \times TVu-7778 population, a framework consensus map was generated by using five populations. The Dan IIa \times TVu-7778 population was added while making sure that no LG or marker order conflicts were introduced.

Within-LG marker assignment was attempted first by using JoinMap; however, major reshuffling of marker order was observed compared with individual maps. Previous studies have encountered the same limitation when JoinMap was used to construct high-density consensus maps (12, 13). Therefore, in our mapping protocol, homologous LGs were used to generate consensus LGs one at a time with MergeMap (12) to establish marker order (see *Materials and Methods*). Then map distances were added by using the cM values generated by JoinMap for each LG. The resulting consensus map contained 928 SNP markers on 619 unique map positions distributed over 11 LGs, covering

Table 2. Pairwise comparison of common SNP markers across cowpea RIL populations

RIL population	524B \times IT84S-2049	CB46 \times IT93K-503-1	TVu-14676 \times IT84S-2246-4	CB27 \times 24-125B-1	Dan Ila \times TVu-7778
Yacine \times 58-77	168	162	161	119	123
Dan Ila \times TVu-7778	147	130	99	130	
CB27 \times 24-125B-1	156	154	108		
TVu14676 \times IT84S-2246-4	137	134			
CB46 \times IT93K-503-1	202				

Table 3. Population size, SNP polymorphisms, map characteristics, and traits for RIL populations used for consensus mapping

Population	RILs	Polymorphic SNPs	SNPs with MAF < 0.3	Mapped SNPs	Map size, cM	Largest gap between SNPs, cM	Traits segregating
524B × IT845-2049	79	469	33	436	665	13.3	AR, IGW, NR, SR, VR, SCT
CB27 × 24-125B-1	90	366	67	299	651	17.0	AR, IGW, CWR, NR, M, FLTR,
CB46 × IT93K-503-1	103	422	34	388	601	17.1	NR, IGW, DT, FLTR, M, MR, Y, AR, FR
Dan IIa × TVu-7778	109	524	236	288	665	16.3	DT, Y, BB
TVu-14676 × IT845-2246-4	137	409	60	349	600	17.9	SR, NR
Yacine × 58-77	114	499	84	415	657	17.6	FLTR, IGW,

AR, aphid resistance; BB, bacterial blight resistance; CWR, cowpea weevil resistance; DT, drought tolerance; FR, Fusarium resistance; FLTR, flower thrips resistance; IGW, individual grain weight; M, maturity; MR, *Macrophomina* resistance; NR, nematode resistance; SR, *Striga* resistance; SCT, seedling cold tolerance; VR, virus resistance; Y, yield.

a total genetic distance of 680 cM. This is an average marker distance of 0.73 cM, or one SNP per 668 kbp considering the cowpea genome to be 620 Mbp. Coincidentally, the map contains almost exactly an average of 1.0 map positions per Mbp (619 map positions/620 Mbp). The average distance between unique map positions was 1.09 cM (680 cM/619 map positions).

Marker density was generally consistent throughout the map with only 19 instances where distances between adjacent markers were >4 cM and only one region on VuLG1 where the distance was >10 cM. VuLG1 also had the least marker density with an average distance of 1.23 cM, whereas VuLG3 had the highest marker density with an average distance of 0.49 cM between markers. The remaining LGs had average marker distances <1.0 ranging from 0.60 to 0.98 cM. LG sizes ranged from 44.75 cM for the smallest to 85.24 cM for the largest. Number of markers, average marker distance, and corresponding LGs from previous publications are summarized in Table 4 and marker distribution across LGs is shown in Fig. S1.

Of the 928 SNP-harboring cowpea unigenes, 921 were annotated based on sequence homology with soybean at e-scores of 1.00e-10 or better. The annotation information for the mapped SNPs is available from HarvEST: Cowpea (<http://harvest.ucr.edu> and www.harvest-web.org). The HarvEST BLAST server (<http://138.23.191.145/blast/index.html>) provides the mapped SNP unigenes as a searchable database.

Synteny with Soybean. Extensive macrosynteny and microsynteny were observed between cowpea and soybean. Only 7 of the 928 genes placed on the cowpea consensus map did not have a soybean hit of e-10 or better. Of the remaining 921 genes, 789 were in regions highly syntenic and collinear with soybean chromosomal (GmChr) regions, which represents ≈85% of the cowpea genome covered by the current map. The ranked order of syntenic regions of soybean is included in Table 4. The number of major soybean synteny blocks for each cowpea LG ranged from one to three, whereas the total

number of significant soybean synteny blocks ranged from three to six. Seven of 11 LGs had major synteny with two soybean chromosomes, among which four (VuLG2, VuLG3, VuLG5, and VuLG7) had synteny along the entire LG. For example, VuLG5 was completely syntenic with soybean 14 as shown in Fig. 1. An additional example is shown in Fig. S2.

Synteny with *M. truncatula*. As expected, synteny was reduced in *M. truncatula* compared with soybean. However, regions with extensive macrosynteny, microsynteny, and collinearity were observed. Of the 928 EST-derived SNPs mapped in the cowpea consensus map, 809 had significant *M. truncatula* hits, with 759 (82%) in regions defined by synteny. Extensive chromosomal rearrangement was evident, with cowpea LGs typically chimeric to three or more *M. truncatula* chromosomal segments (Table 4). Regardless, blocks of extensive synteny and collinearity were evident, because 10 of the 11 cowpea LGs had at least 17 corresponding loci on a *Medicago* chromosome (Table 4). Specifically, the entire VuLG7 exhibited extensive synteny and collinearity with a section of MtChr5 (Fig. 2). Similarly, VuLG2 had extensive macrosynteny with MtChr1 where 76 of 116 cowpea loci had corresponding orthologs on MtChr1 (Table 4). Also, one block of VuLG3 was syntenic and collinear with the lower section of MtChr4.

Synteny with *Arabidopsis*. Major chromosomal rearrangement was observed between cowpea and *Arabidopsis* such that no macrosynteny was evident. Significant microsynteny was observed in some regions, but collinearity was markedly reduced relative to legume reference genomes. The strongest instance of cowpea-*Arabidopsis* microsynteny and collinearity was an ≈14-cM section of VuLG1 and AtChr1 where gene order exhibited only minor differences.

Discussion

In this study, the high-throughput Illumina GoldenGate SNP assay was tested and validated for cowpea by using six RIL mapping populations.

Table 4. Consensus genetic map, correspondence of LGs to previous maps, and synteny with soybean (GmChr) and *Medicago* (MtChr) highlighted by number of cowpea orthologs in parentheses

Consensus VuLG	Markers	Length	Ouédraogo et al. (6) LG	Muchero et al. (8) LG	GmChr (Cowpea orthologs)	MtChr (Cowpea orthologs)
1	69	85.2	4	6	18(27), 9(10), 13(9), 8(8), 15(5)	7(25), 2(12), 3(7), 4(7)
2	116	84.0	4,11	5, 9	10(43), 20(39), 2(13)	1(76), 7(15), 4(12), 5(11), 2(5)
3	168	81.8	3	5,10	5(43), 8(39), 17(33), 7(15), 13(14), 16(5)	4(41), 8(36), 5(23), 2(20), 3(12), 6(6)
4	68	66.4	9	7, 8	19(19), 3(13), 11(9), 18(9), 16(5)	3(22), 7(22), 5(5)
5	75	62.6	5	11	14(37), 2(15), 17(9)	5(38), 1(14), 3(9)
6	93	59.1	8	3	15(32), 8(15), 9(10), 13(8), 18(9), 19(6)	2(37), 3(11), 5(11), 7(7), 1(6), 4(6), 8(6)
7	72	52.9	5	1	11(21), 1(20), 2(9), 9(9)	5(42), 1(6), 8(6), 7(5)
8	65	49.0	7,10	2	6(27), 4(26), 15(5)	3(29), 8(9), 4(7)
9	66	48.4	1	1	12(30), 11(11), 15(9), 6(7)	4(17), 2(15), 8(8), 3(7)
10	77	46.0	6	2, 3	7(23), 3(20), 1(10), 16(10), 8(6)	8(21), 4(14), 7(12), 5(8), 2(7)
11	59	44.8	2	2, 3	16(15), 13(10), 19(9), 9(8), 2(5)	6(12), 4(11), 5(6), 1(5), 7(5)




Fig. 1. Macrosynteny and collinearity between cowpea LG VuLG5 and soybean chromosomes GmChr2, GmChr14, and GmChr17.




Fig. 2. Macrosynteny and collinearity between cowpea linkage VuLG7 and *M. truncatula* chromosomes MtChr6.

The effectiveness and utility of this approach in cowpea was demonstrated by the successful genotyping of 1,375 of 1,536 SNP loci (89.55%) among 741 RILs derived from six crosses, synthetic heterozygotes, and parental genotypes. Further, the quality of the data was highlighted by the incorporation of 928 (67%) of these 1,375 markers into a consensus genetic map. Given that this oligonucleotide pool assay was developed for cowpea and none of the SNPs were validated in practice before this work to our knowledge, the 90% success rate supports the efficacy of strategies adopted in developing EST-derived SNPs from cowpea.

The gene-based consensus map described here will enable integration of information from populations developed from diverse crosses, enable whole genome molecular marker-assisted selection, and facilitate linkage disequilibrium studies, association mapping, and synteny-based genomics in cowpea. All of these possibilities should benefit from the dense marker placement (0.73 cM/marker) in the current map. Significantly, almost all syntenic blocks were in subtelomeric regions of soybean chromosomes, suggesting that these are gene-rich and that the heterochromatic pericentric regions are not densely populated with markers in the cowpea map. Although LGs <50 cM for VuLG8, VuLG9, VuLG10, and VuLG11 might suggest incomplete maps, this observation could result from high within-LG marker ordering LODs. All four LGs had lengths >50 cM using LODs between 1 and 3. To minimize spurious linkages, we used LODs 7 or higher. The map size reported here was consistent with previously published map sizes of 972 cM (7), 643 cM (8), and 669.8 cM (9). However, the Ouédraogo et al.

(6) map was ≈ 3 –4 times longer with a total distance of 2,670 cM. The same mapping population (524B \times IT84S-2049) used to construct that map was used in the current study, and its individual SNP-based map size was 665 cM, well within the expected range, based on the average observed across the individual maps.

EST-derived SNP markers provide an additional level of utility in genomic studies because they tag actual genes that, when incorporated into genetic maps, may be used for synteny-based cross-species genome comparison. Soybean genome duplication was evident because a single cowpea haplotype was typically syntenic with regions on at least two different soybean chromosomes. Further, chromosomal rearrangement was apparent between the two species, because each cowpea LG was syntenic to regions on at least three and up to six soybean chromosomes. Although genome duplication in soybean will present a challenge in translational genomics with cowpea, extensive macrosynteny and collinearity in duplicated regions compared with cowpea should facilitate such efforts. The observation that 789 of the 928 EST-derived SNP markers mapped in regions exhibiting synteny and conserved collinearity suggests that $\approx 85\%$ of the cowpea genome covered by this map has strong correspondence to at least one soybean genomic region. Therefore, studies in cowpea should benefit from the extensive soybean genome sequence resource in identifying genetic determinants of traits of economic importance. Similarly, because cowpea is one of the most productive crops in hot, low rainfall, and poor soil environments (2, 4), studies in soybean and common bean should benefit from validation of

ACKNOWLEDGMENTS. We thank E. Lindquist, C. Pennacchio, S. Lucas, M. Wang, and J. Bristow of the U.S. Department of Energy Joint Genome Institute (Walnut Creek, CA) for EST sequencing of libraries UCRVU04 through UCRVU12; M. Timko (University of Virginia, Charlottesville) for providing seeds of the Dan Ila \times TVu-7778 and IT84S-2246 \times TVu-14676 RIL populations; C. Sudhakar (Sri Krishnadevaraya University, Anantapur, India) for production of one of the cDNA libraries; and A. Farmer (National

Center for Genome Resources, Santa Fe, NM) for a list of soybean gene models related to a National Science Foundation project led by D. Cook (University of California, Davis). This work was supported by the Generation Challenge Program through a grant from the Bill and Melinda Gates Foundation and U.S. Agency for International Development Collaborative Research Support Program Grants GDG-G-00-02-00012-00 and EDH-A-00-07-00005.

1. Delmer DP (2005) Agriculture in the developing world: Connecting innovations in plant research to downstream applications. *Proc Natl Acad Sci USA* 102:15739–15746.
2. Ehlers JD, Hall AE (1997) Cowpea (*Vigna unguiculata* L. Walp). *Field Crops Res* 53:187–204.
3. Sanginga N, et al. (2003) Sustainable resource management coupled to resilient germplasm to provide new intensive cereal–grain–legume–livestock systems in the dry savanna. *Agric Ecosys Environ* 100:305–314.
4. Hall AE (2004) Breeding for adaptation to drought and heat in cowpea. *Eur J Agron* 21:447–454.
5. Timko M, et al. (2008) Sequencing and analysis of the gene-rich space of cowpea. *BMC Genomics* 9:103.
6. Ouédraogo JT, et al. (2002) An improved genetic linkage map for cowpea (*Vigna unguiculata* L.) Combining AFLP, RFLP, RAPD, biochemical markers, and biological resistance traits. *Genome* 45:175–188.
7. Menéndez CM, Hall AE, Gepts P (1997) A genetic linkage map of cowpea (*Vigna unguiculata*) developed from a cross between two inbred, domesticated lines. *Theor Appl Genet* 95:1210–1217.
8. Muchero W, Ehlers J, Close T, Roberts P (2009) Mapping QTL for drought stress-induced premature senescence and maturity in cowpea [*Vigna unguiculata* (L.) Walp.]. *Theor Appl Genet* 118:849–863.
9. Ubi BE, Mignouna H, Thottappilly G (2000) Construction of a genetic linkage map and QTL analysis using a recombinant inbred population derived from an intersubspecific cross of cowpea [*Vigna unguiculata* (L.) Walp.]. *Breed Sci* 50:161–172.
10. Young ND, Mudge J, Ellis TN (2003) Legume genomes: More than peas in a pod. *Curr Opin Plant Biol* 6:199–204.
11. Van Ooijen JW, Voorrips RE (2001) *JoinMap 3.0: Software for the Calculation of Genetic Linkage Maps* (Plant Research International, Wageningen, The Netherlands).
12. Wu Y, Close TJ, Lonardi S (2008) in *Computational Systems Bioinformatics: CSB 2008 Conference*, eds Markstein P, Xu Y (Imperial College Press, Stanford, CA), pp 285–296.
13. Yap IV, et al. (2003) A graph-theoretical approach to comparing and integrating genetic, physical, and sequence-based maps. *Genetics* 165:2235–2247.
14. Huang X, Madan A (1999) CAP3: A DNA sequence assembly program. *Genome Res* 9:868–877.
15. Ismail AM, Hall AE, Close TJ (1999) Allelic variation of a dehydrin gene cosegregates with chilling tolerance during seedling emergence. *Proc Natl Acad Sci USA* 96:13566–13570.
16. Mould MJR, Xu T, Barbara M, Iscove NN, Heath MLC (2003) cDNAs generated from individual epidermal cells reveal that differential gene expression predicting subsequent resistance or susceptibility to rust fungal infection occurs prior to the fungus entering the cell lumen. *Mol Plant-Microbe Interact* 16:835–845.
17. Simões-Araújo JL, et al. (2002) Identification of differentially expressed genes by cDNA-AFLP technique during heat stress in cowpea nodules. *FEBS Lett* 515:44–50.
18. Kosambi DD (1944) The estimation of map distance from recombination values. *Ann Eugen* 12:172–175.
19. Voorrips RE (2002) MapChart: Software for the graphical presentation of linkage maps and QTLs. *J Heredity* 93:77–78.